

Systematic Evaluation of the Dependence of Deoxyribozyme Catalysis on Random Region Length


Tania E. Velez,[†] Jaydeep Singh,[†] Ying Xiao, Emily C. Allen, On Yi Wong, Madhavaiah Chandra, Sarah C. Kwon, and Scott K. Silverman*

Department of Chemistry, University of Illinois at Urbana–Champaign, 600 South Mathews Avenue, Urbana, Illinois 61801, United States

Supporting Information

ABSTRACT: Functional nucleic acids are DNA and RNA aptamers that bind targets, or they are deoxyribozymes and ribozymes that have catalytic activity. These functional DNA and RNA sequences can be identified from random-sequence pools by in vitro selection, which requires choosing the length of the random region. Shorter random regions allow more complete coverage of sequence space but may not permit the structural complexity necessary for binding or catalysis. In contrast, longer random regions are sampled incompletely but may allow adoption of more complicated structures that enable function. In this study, we systematically examined random region length (N_{20} through N_{60}) for two particular deoxyribozyme catalytic activities, DNA cleavage and tyrosine-RNA nucleopeptide linkage formation. For both activities, we previously identified deoxyribozymes using only N_{40} regions. In the case of DNA cleavage, here we found that shorter N_{20} and N_{30} regions allowed robust catalytic function, either by DNA hydrolysis or by DNA deglycosylation and strand scission via β -elimination, whereas longer N_{50} and N_{60} regions did not lead to catalytically active DNA sequences. Follow-up selections with N_{20} , N_{30} , and N_{40} regions revealed an interesting interplay of metal ion cofactors and random region length. Separately, for Tyr-RNA linkage formation, N_{30} and N_{60} regions provided catalytically active sequences, whereas N_{20} was unsuccessful, and the N_{40} deoxyribozymes were functionally superior (in terms of rate and yield) to N_{30} and N_{60} . Collectively, the results indicate that with future in vitro selection experiments for DNA and RNA catalysts, and by extension for aptamers, random region length should be an important experimental variable.

KEYWORDS: deoxyribozyme catalysis, nucleic acids, DNA catalysts, RNA catalysts, random region length, DNA and RNA aptamers



INTRODUCTION

Naturally occurring RNA aptamers form key binding elements of riboswitches,¹ and natural RNA enzymes (ribozymes) catalyze RNA cleavage and ligation reactions² as well as peptide bond formation in the ribosome.³ In contrast, natural DNA aptamers or enzymes (deoxyribozymes) have not been reported. The identification of the first natural ribozymes spurred searches for artificial aptamers and nucleic acid enzymes, comprising both RNA^{4–6} and soon thereafter DNA.^{7–9} Such searches are performed using in vitro selection, which is a powerful approach for identifying new, functional DNA or RNA sequences from random-sequence “pools”.^{10,11} Many artificial DNA and RNA aptamers¹² and enzymes^{13–16} have been found using selection methodologies.

In nearly all selection experiments, the length n of the pool's random region is fixed at the outset of the experiment. With only four standard nucleotide monomers, the size of sequence space increases as 4^n . The physical sample of the initial pool is typically prepared by solid-phase synthesis for a DNA pool, or by transcription using a DNA template itself prepared by solid-phase synthesis for an RNA pool. As a practical limitation, the amount of the starting pool is on the order of 10^{14} molecules (200 pmol), or perhaps 10^{15} or 10^{16} in some cases. In

comparison, the 4^n size of sequence space can be considerably larger; for example, for a 60-nucleotide N_{60} pool, $4^{60} \approx 10^{36}$. The sampling of sequence space is extremely sparse for long random regions. For the N_{60} case, $10^{14}/10^{36} = 10^{-22}$ is the fraction of sequence space sampled. To the best of our knowledge, the longest reported random region is N_{228} , for which only 10^{-123} of sequence space is covered.¹⁷ Toward the other end of the length spectrum, an N_{20} region has only $4^{20} = 10^{12}$ possible sequence permutations, and a typical starting pool with 10^{14} molecules will sample essentially every possible 20-nucleotide sequence (10^2 molecules of each N_{20} sequence, on average).

In any particular selection experiment, the choice of random region length n requires a compromise between two competing considerations: (1) coverage of sequence space, where shorter random regions allow greater coverage; and (2) potential structural complexity of the nucleic acid motifs, where longer random regions allow greater complexity of secondary and tertiary structure. Another consideration is that longer random

Received: September 12, 2012

Revised: October 16, 2012

Published: October 22, 2012

regions may suffer from increased propensity to misfold, or from inhibition of catalytic portions by the excess sequence elements.¹⁸ The choice of n is typically made in a relatively arbitrary fashion; some laboratories may prefer N_{40} pools, others N_{50} , and others N_{70} , and each lab rarely if ever uses one of the “other” lengths. Without experimentally assessing the emergent binding or catalysis as a systematic function of random region length, one cannot be confident in choosing the best compromise between sequence space coverage and structural complexity.

Theoretical calculations have explored the probabilities of finding various structural motifs in random-sequence pools,^{18–20} among other sophisticated computational analyses of selection processes.^{21–25} However, experiments are the ultimate arbiter of success in identifying ligands or catalysts. Only a small number of reports have strived to examine empirically the importance of random region length for a particular binding or catalytic activity. Huang et al. sought ribozymes for coenzyme synthesis by mixing pools of widely varying lengths (N_{30} , N_{60} , N_{100} , and N_{140}), finding activity only from the N_{30} and N_{60} sequences.^{26,27} However, under these competitive selection conditions, an additional consideration is the strong replicative advantage of shorter sequences, which is not an issue when different length pools are studied separately rather than competing directly.²⁸ Yarus and co-workers examined random regions from length 16 to 90 under noncompetitive conditions to identify RNA aptamers for the amino acid isoleucine, finding that 50 or 70 nt random regions were optimal.²⁹

We are unaware of studies in which any kind of deoxyribozyme catalysis has been examined as a function of random region length. In the present study, we chose two particular catalytic activities for systematic analysis (Figure 1). First, we examined DNA-catalyzed DNA cleavage, where we have previously identified deoxyribozymes with rate enhancements $>10^{13}$ for phosphodiester bond hydrolysis.^{30,31} Second, we examined DNA-catalyzed nucleopeptide linkage formation, which we have identified as part of our larger efforts that seek DNA-catalyzed covalent modification of amino acid side chains.^{32–37} For both types of catalytic activity, our previous efforts used only N_{40} random regions, where $4^{40} \approx 10^{24}$ and therefore $10^{14}/10^{24} = 10^{-10}$ of sequence space was explored. Here, for both kinds of catalysis we examined N_{20} through N_{60} pools in new, independent selection experiments. For N_{20} , sequence space is fully covered, as noted above. For N_{30} , N_{50} , and N_{60} , the fractions of covered sequence space are approximately 10^{-4} , 10^{-16} , and 10^{-22} , respectively. For both DNA-catalyzed reactions, the results showed that the length of the random region substantially impacts the outcome of the selection process, with important implications for future in vitro selection experiments.

RESULTS AND DISCUSSION

DNA-Catalyzed DNA Cleavage: Selection Experiments with N_{20} through N_{60} Random Regions. Our previous in vitro selection experiments to identify deoxyribozymes for DNA cleavage have used solely N_{40} random regions.^{30,31,38–40} Here, we separately performed analogous experiments using N_{20} , N_{30} , N_{50} , and N_{60} pools. The selection process was performed using our previously described approach,^{31,41,42} with a single-stranded DNA substrate and a deoxyribozyme that interacts with the substrate via two fixed Watson–Crick binding arms (Figure 1A). Each iterated selection round




Figure 1. Two reactions systematically examined for the effect of random region length on DNA catalysis. In each case, the key selection step is illustrated. (A) DNA-catalyzed DNA cleavage. (B) DNA-catalyzed tyrosine-RNA nucleopeptide linkage formation.

consisted of three steps of (1) PCR amplification of the population from the previous round, (2) enzymatic ligation of the DNA substrate to the amplified pool, and finally (3) incubation to allow individual active deoxyribozyme sequences to cleave their attached DNA substrate. Active deoxyribozyme sequences were separated by polyacrylamide gel electrophoresis (PAGE) on the basis of the downward shift due to loss of approximately half of the DNA substrate. During this key selection step, the incubation conditions were 70 mM HEPES, pH 7.5, 1 mM $ZnCl_2$, 20 mM $MnCl_2$, 40 mM $MgCl_2$, and 150 mM NaCl at 37 °C for 14 h. The resulting deoxyribozymes function without covalent attachment to the DNA substrate via the loop that is remote from the hydrolysis site.

After nine selection rounds, the N_{50} and N_{60} experiments showed no detectable activity ($<0.2\%$) and were discontinued. In contrast, the N_{20} and N_{30} selections led to detectable ($>0.2\%$) cleavage activity beginning at round 4 (N_{20} , 0.9%) and round 5 (N_{30} , 7.5%). Each pool was cloned from round 8 (48% and 45% for N_{20} and N_{30} , respectively), and individual deoxyribozymes were characterized (see sequences in Supporting Information, Figure S1).

From the N_{20} pool, nine individual deoxyribozymes were obtained, exhibiting a range of cleavage sites along the DNA substrate (Figure 2A). The cleavage rate constant k_{obs} was as high as 0.2 h^{-1} (Figure 3A and Supporting Information, Figure S2A). MALDI mass spectrometry revealed that two of these nine N_{20} deoxyribozymes catalyze site-specific DNA phosphodiester hydrolysis, whereas seven catalyze deglycosylation followed by strand scission via two β -elimination reactions (Figure 4 and Supporting Information, Tables S1 and S2). Both we⁴⁰ and others⁴³ have observed deglycosylation-induced DNA




Figure 2. Various substrate cleavage sites for deoxyribozymes from the initial (A) N_{20} and (B) N_{30} selections for DNA-catalyzed DNA cleavage. Individual deoxyribozymes are denoted 8VA followed by a clone number for N_{20} and 8VB followed by a clone number for N_{30} . Each deoxyribozyme was allowed to cleave the 5'- 32 P-radiolabeled DNA substrate (*open arrowhead*), resulting in a product band (*filled arrowhead*) whose PAGE migration rate relative to the standard suggests the cleavage site. The standard sample corresponds to hydrolysis of the substrate sequence at the position noted in the Experimental Procedures. Incubation conditions: 70 mM HEPES, pH 7.5, 1 mM $ZnCl_2$, 20 mM $MnCl_2$, 40 mM $MgCl_2$, and 150 mM NaCl at 37 °C ($t = 20$ h). Quantification of cleavage kinetics for each deoxyribozyme is provided in Figure 3. The reaction type (i.e., substrate hydrolysis, or substrate deglycosylation followed by two β -eliminations) and precise site of each cleavage reaction was assigned with confidence for each deoxyribozyme by MALDI mass spectrometry of the products (Figure 4; see Supporting Information, Tables S1 and S2 for all data values and cleavage-site assignments). The type of reaction catalyzed by each deoxyribozyme is marked below its lane with "H" for hydrolysis or "D" for deglycosylation.




Figure 3. Kinetic plots for individual deoxyribozymes from the (A) N_{20} and (B) N_{30} selections for DNA cleavage. Kinetic plots for several additional deoxyribozymes from each selection experiment are shown in Supporting Information, Figure S2. k_{obs} values are tabulated in Supporting Information, Table S4. Incubation conditions as in Figure 2.

cleavage by unrelated deoxyribozymes. Similar results were found for the N_{30} pool, which led to 14 individual deoxyribozymes with a range of cleavage sites (Figure 2B). The cleavage k_{obs} was as high as 0.5 h^{-1} (Figure 3B and Supporting Information, Figure S2B). MALDI mass spectrom-

etry indicated that three of these N_{30} deoxyribozymes hydrolyze the DNA substrate site-specifically, whereas 11 induce deglycosylation and strand scission (Figure 4 and Supporting Information, Tables S1 and S2). Curiously, our selection experiments with N_{40} random regions led almost entirely to hydrolysis,³¹ whereas the N_{20} and N_{30} selections described here led to many deoxyribozymes that catalyze deglycosylation and strand scission via β -elimination. This unexpected functional distinction between N_{20}/N_{30} and N_{40} selections highlights the importance of random region length as a variable that should be explored during selection experiments.

DNA-Catalyzed DNA Hydrolysis: Follow-Up Experiments to Explore Interplay between Metal Ion Cofactors and Random Region Length. From the above-described efforts along with our previous work,^{30,31,38–40} we inferred that the longer random-region lengths (N_{50} and N_{60}) are sampled too sparsely for successful identification of deoxyribozymes that catalyze DNA cleavage, whereas N_{20} , N_{30} , and N_{40} permit such identification. Excess sequence elements in the N_{50} and N_{60} sequences may also have suppressed the function of what otherwise would be active catalysts derived from shorter DNA segments. Whichever explanation applies, the outcome is that no activity was observed from the N_{50} and N_{60} pools, but the N_{20} and N_{30} (and N_{40}) pools led to substantial catalytic function. We noted that the majority of the new N_{20} and N_{30} deoxyribozymes cleave their DNA substrate by a nonhydrolytic pathway, whereas hydrolysis is the more desirable outcome, for example, for preparative purposes. We also considered that all of the prior and current selection experiments were performed with both Zn^{2+} and Mn^{2+} (as well as Mg^{2+}) as available metal ion cofactors. In many cases, the outcome was that both Zn^{2+} and Mn^{2+} were required for catalysis, although one of our earlier (N_{40}) DNA-hydrolyzing deoxyribozymes was subsequently shown to require only Zn^{2+} ,³⁹ as were several deoxyribozymes identified later.⁴⁰

We therefore performed a set of follow-up selection experiments, with two key changes to the procedure. First, we implemented a "capture step" immediately after each selection step, using T4 DNA ligase and a splinted 5'-




Figure 4. Representative MALDI mass spectra to assign cleavage reactions and sites for deoxyribozymes from the initial N_{20} and N_{30} selections for DNA cleavage. All mass spectra data and precise cleavage-site assignments are tabulated in Supporting Information, Tables S1 and S2. The indicated C nucleotide of the substrate was not base-paired with either deoxyribozyme binding arm (see Figure 1). (A) 8VB4 deoxyribozyme, which hydrolyzes the DNA substrate at a specific phosphodiester linkage and forms 3'-phosphate and 5'-hydroxyl products. Some of the new DNA-hydrolyzing deoxyribozymes lead instead to 3'-hydroxyl and 5'-phosphate products. (B) 8VB5 deoxyribozyme, which deglycosylates the DNA substrate at a specific guanosine. After two subsequent β -elimination reactions, the products are missing the entire G nucleoside and have 3'-phosphate and 5'-phosphate groups. Some of the new DNA-deglycosylating deoxyribozymes are not site-specific and deglycosylate the substrate at either of two adjacent nucleotide positions. In the spectrum of panel B, the asterisk denotes the peak for uncleaved substrate with $z = 2$.

phosphorylated donor oligonucleotide to capture the 3'-hydroxyl group uniquely revealed by DNA-catalyzed DNA hydrolysis (see Experimental Procedures) but not by deglycosylation, which after strand scission leads to a 3'-phosphate group. Selection pressure via a similar capture step was implemented in one of our previous reports (see Figure 5 in ref 31, which depicts capture of the 5'-phosphate product as the selection pressure to achieve cleavage-site specificity). Second, in addition to including both Zn^{2+}/Mn^{2+} during the selection step, we separately evaluated providing Zn^{2+} alone as the divalent metal ion cofactor. Because we have already reported the outcome from evaluating N_{40} with Zn^{2+}/Mn^{2+} ,³¹ here we performed five new follow-up selection experiments: N_{20} and N_{30} with Zn^{2+}/Mn^{2+} , and N_{20} , N_{30} , and N_{40} with Zn^{2+} alone, all five now with inclusion of selection pressure via the capture step to enforce DNA-catalyzed DNA hydrolysis.

Of these five follow-up selection experiments, only the combination of N_{30} with Zn^{2+} and Mn^{2+} led to no activity through 12 selection rounds; this experiment was discontinued. The other four efforts all led to considerable catalytic activity, and each was cloned (N_{20} with Zn^{2+} : 26% at round 7; N_{20} with Zn^{2+}/Mn^{2+} : 4% at round 9; N_{30} with Zn^{2+} , 25% at round 7; N_{40} with Zn^{2+} , 12% at round 9). Each experiment led to just one or two distinct deoxyribozyme sequences (Supporting Information, Figure S1), with k_{obs} as high as 0.5 h^{-1} (Figure 5). MALDI mass spectrometry confirmed that all of these deoxyribozymes catalyze site-specific DNA hydrolysis at the expected location within the DNA substrate, with formation of 3'-hydroxyl and 5'-phosphate termini (Supporting Information, Table S1). Nonhydrolytic cleavage via deglycosylation and strand scission was not observed by any of these deoxyribozymes, consistent with the stringent nature of the applied selection pressure. Notably, all five (out of five) of the N_{20} and N_{30} deoxyribozymes from these follow-up selection experiments led to DNA hydrolysis, whereas in the first set of N_{20} and N_{30} selections, only five out of 23 did so; clearly the selection pressure via the splint ligation capture step is highly effective.

All four of the new $N_{20}/N_{30}/N_{40}$ DNA-hydrolyzing deoxyribozymes that were identified with Zn^{2+} alone were tested briefly for substrate sequence generality, by varying all DNA substrate nucleotides except for the unpaired C[^]G at the substrate cleavage site and covarying the deoxyribozyme




Figure 5. Activities of deoxyribozymes from the follow-up N_{20} , N_{30} , and N_{40} selections for DNA-catalyzed DNA hydrolysis. (A) PAGE image for each of the six new deoxyribozymes, showing representative time points ($t = 30 \text{ s}$, 15 min , 2 h , and 20 h). Substrate (open arrowhead) and product (filled arrowhead) are marked. Incubation conditions: 70 mM HEPES, $\text{pH } 7.5$, 1 mM $ZnCl_2$, 20 mM $MnCl_2$ if indicated, and 150 mM $NaCl$ at $37 \text{ }^\circ\text{C}$. (B) Kinetic plots. k_{obs} values are tabulated in Supporting Information, Table S4.

binding arms to maintain Watson–Crick base pairing.³¹ None of these four deoxyribozymes retained detectable ($>0.2\%$) catalytic activity when the substrate nucleotides were changed systematically in this fashion ($A \leftrightarrow T$, $G \leftrightarrow C$; data not shown). Because our previous N_{40} experiments with Zn^{2+}/Mn^{2+} did lead to numerous highly general DNA-hydrolyzing deoxyribozymes,³¹ the present finding indicates that the use of Zn^{2+} alone as the cofactor comes at an apparent functional cost, namely, the loss of tolerance for a range of DNA substrate

sequences. We previously observed a similar compromise for our lone prior example of an N_{40} deoxyribozyme that is active with Zn^{2+} alone.³⁸

We did not anticipate the experimental outcome that the shorter N_{20} and N_{30} pools lead to better catalysis with Zn^{2+} alone rather than with Zn^{2+} and Mn^{2+} . In Figure 5, note that 9ZHS—identified with N_{20} and Zn^{2+}/Mn^{2+} —had the worst activity of all six new deoxyribozymes, and the N_{30} selection with Zn^{2+}/Mn^{2+} led to no activity at all. This finding highlights the relatively “rugged” nature of selection landscapes, in that changing the pool length from N_{40} to N_{20}/N_{30} unpredictably changed the optimal metal ions from Zn^{2+} and Mn^{2+} together to Zn^{2+} alone. These results establish an interesting interplay for DNA-catalyzed DNA hydrolysis between two key selection variables, random region length and divalent metal ion cofactors. A more general implication of these results is that one must be careful about assuming that a particular finding (e.g., with DNA hydrolysis and an N_{40} pool, Zn^{2+}/Mn^{2+} is superior to Zn^{2+} alone) will apply for different pool lengths (such as N_{20} and N_{30}).

DNA-Catalyzed Nucleopeptide Linkage Formation: Selection Experiments with N_{20} – N_{60} Random Regions.

Separately from the above-described experiments with DNA-catalyzed DNA cleavage, we examined the influence of random region length on DNA-catalyzed nucleopeptide linkage formation, where all of our previous efforts with this reaction have been restricted to N_{40} random regions.^{32–36} New selection experiments with N_{20} , N_{30} , and N_{60} pools were performed as done previously,³⁴ using an “open” architecture and a Cys-Tyr-Ala tripeptide substrate connected via a flexible hexa(ethylene glycol) (HEG) tether to a DNA anchor oligonucleotide (Figure 1B). The electrophile for reaction with the tyrosine nucleophile was a 5'-triphosphorylated RNA strand. Iterated selection rounds were performed with incubation in 50 mM HEPES, pH 7.5, 20 mM $MnCl_2$, 40 mM $MgCl_2$, 150 mM NaCl, and 2 mM KCl at 37 °C for 2 h in each key selection step. Active deoxyribozyme sequences were separated on the basis of their upward PAGE shift due to attachment of the RNA strand to the DNA-anchored Cys-Tyr-Ala tripeptide.

The impact of the random region length on the outcome of DNA-catalyzed nucleopeptide linkage formation was found to be very different from the observations with DNA-catalyzed DNA cleavage as described above. Here, the N_{20} pool led to no detectable activity after nine rounds and was discontinued. In contrast, both of the N_{30} and N_{60} pools led to substantial catalytic activity: N_{30} 35% at round 8 and N_{60} 26% at round 7. Each of these selection experiments was cloned, and individual deoxyribozymes were characterized further. From the N_{30} selection, a single distinct deoxyribozyme sequence was identified (Supporting Information, Figure S1), with rate constant and yield substantially lower than those of the analogous N_{40} deoxyribozymes that were identified previously (k_{obs} 0.15 h^{-1} and 52% yield at 20 h for N_{30} , versus 0.95 h^{-1} and 73% yield for N_{40} ; Figure 6).³⁴ From the N_{60} selection, two deoxyribozymes sequences were found (Supporting Information, Figure S1), and their rate constants and yields were also considerably lower than for N_{40} (k_{obs} 0.07 h^{-1} and 19% yield for the more active variant; Figure 6). MALDI mass spectrometry confirmed that for all of these deoxyribozymes, the product was consistent with formation of the expected Tyr-RNA nucleopeptide linkage (Supporting Information, Table S3). Therefore, for DNA-catalyzed nucleopeptide linkage formation, the outcomes with random region lengths other than N_{40} were markedly




Figure 6. Activities of deoxyribozymes from the selections for DNA-catalyzed nucleopeptide linkage formation. (A) PAGE image showing representative time points ($t = 30$ s, 30 min, 2 h, and 20 h). Incubation conditions: 50 mM HEPES, pH 7.5, 20 mM $MnCl_2$, 40 mM $MgCl_2$, 150 mM NaCl, and 2 mM KCl at 37 °C. Substrate (open arrowhead) and product (filled arrowhead) are marked. For N_{30} , 8TM3 was the only new deoxyribozyme identified, along with two closely related sequence variants. For N_{40} , 11MN5 was the best deoxyribozyme identified previously by direct selection;³¹ data shown here were newly acquired for comparison. For N_{60} , 7TQ20 was one of two different deoxyribozymes found, along with six close sequence variants; the other deoxyribozyme, 7TQ46 (no variants found), had slightly lower activity. (B) Kinetic plots. k_{obs} values are tabulated in Supporting Information, Table S4.

different than the outcomes for DNA-catalyzed DNA hydrolysis, in terms of both (i) which particular random region lengths led to observable catalytic activity, and (ii) quantitative aspects of the catalysis.

General Implications of the Findings for Future In Vitro Selection Experiments. Most laboratories that perform in vitro selection appear to settle on using just one random region length. The specific choice of random region length is usually arbitrary or at least not specifically justified, except perhaps by reference to past successful selections using the same length. Our own laboratory has generally favored N_{40} , but without any systematic examination (until now) of shorter or longer random regions. Similarly, others typically use N_{60} or other lengths. We undertook the present study to provide discrete experimental data regarding impact of random region length on identification of DNA catalysts. The most important implication of our collective results is that one particular random region length is not optimal for all catalytic activities, and therefore the planning of in vitro selection experiments should prominently take into account “random region length”

as an important experimental variable. This is a nontrivial consideration, given the wide range of other variables that may also be adjusted along with the reality that a finite number of selection experiments can be performed in parallel, as well as the practical reality that most reported *in vitro* selection efforts have used only one random region length within each study. We are currently making use of the present findings in several experiments underway in our laboratory, in which we are now systematically evaluating random region lengths other than our previously standard N_{40} . Although the results described here were obtained specifically in the context of DNA catalysts, we anticipate that they also apply to RNA catalysts as well as both DNA and RNA aptamers.

■ EXPERIMENTAL PROCEDURES

Oligonucleotide Preparation. DNA oligonucleotides were prepared by solid-phase synthesis at Integrated DNA Technologies (Coralville, IA) and purified by denaturing 20% or 8% PAGE. To avoid chelation of Zn^{2+} by adventitious EDTA, all substrates and deoxyribozymes used in kinetic assays were extracted from gels using TN buffer (10 mM Tris, pH 8.0, 300 mM NaCl) lacking EDTA and precipitated with ethanol.

In Vitro Selection for DNA-Catalyzed DNA Cleavage.

In vitro selection for DNA-catalyzed DNA cleavage was performed essentially as described previously, using 200 pmol ($\sim 10^{14}$ molecules) in each initial round of selection.^{31,41,42}

For the initial selection experiments (N_{20} , N_{30} , N_{50} , and N_{60} , each with $Zn^{2+}/Mn^{2+}/Mg^{2+}$), the deoxyribozyme pool strand was 5'-CGAAGTCGCCATCTCTTC- N_x -ATAGTGAGTCG-TATTAAGCTGATCCTGATGG-3', where the two underlined regions denote the substrate binding arms. The 5'-CGAA was replaced with 5'-CC and the 3'-terminus was ...TATTA-3' for individual deoxyribozymes prepared by solid-phase synthesis and used in the single-turnover *in trans* kinetic assays. The two PCR primers used during selection were 5'-CGAAGTCGCCATCTCTTC-3' (5'-phosphorylated to enable ligation by T4 RNA ligase) and 5'-(AAC)₄XCCATCAGGATCAGCT-3' (where X denotes Glen Spacer 18, which is a PEG spacer that stops extension by Taq polymerase and leads to a size difference between the two PCR product strands). The DNA substrate used during selection was 5'-TAATACGACTCACTATCGAAGAGATGGCGACgga-3', where the underlined C was unpaired and the three 3'-terminal nucleotides were RNA to enable ligation by T4 RNA ligase to the 5'-terminus of the deoxyribozyme pool strand. The DNA substrate for single-turnover *in trans* assays of individual deoxyribozymes was the all-DNA version of the substrate used during selection. The standard sample used in Figure 2 corresponds to hydrolysis immediately to the 5'-side of the unpaired C, with formation of 3'-hydroxyl and 5'-phosphate termini.

For the follow-up selection experiments (N_{20} and N_{30} each with either Zn^{2+} or Zn^{2+}/Mn^{2+} , or N_{40} with Zn^{2+}), the deoxyribozyme pool strand and individual deoxyribozymes for kinetic assays were the same as in the first set of selection experiments. The two PCR primers used during selection were the same as in the first set of selection experiments, except the first primer was not 5'-phosphorylated. The DNA substrate used during selection and for single-turnover *in trans* kinetic assays of individual deoxyribozymes was 5'-GGATAATACGACTCACTATCGGAAGAGATGGCGACTTCG-3', where the underlined CG was unpaired. The 5'-phosphorylated terminus of the DNA substrate was joined to the 3'-terminus

of the deoxyribozyme pool strand in each round of selection by splint ligation using T4 DNA ligase and a complementary DNA splint (5'-ATAGTGAGTCGATTATCCTCCATCAGGATCAGCTTAATACGACTCACTAT-3'). The capture step in each selection round was performed by splint ligation essentially as described, using T4 DNA ligase.³¹ The splint was 5'-ATCCTGATAACGAGAGGCGATAGTGAGTCG-TATTATCCTCCATCAGGATCAGCT-3', and the 5'-phosphorylated donor (to react with the 3'-OH group revealed after DNA hydrolysis) was 5'-GCCTCTCGTTATCAGGATAACAACAACAACAAC-3'.

In Vitro Selection for DNA-Catalyzed Nucleopeptide Linkage Formation.

In vitro selection for DNA-catalyzed nucleopeptide linkage formation was performed as described previously, using 200 pmol ($\sim 10^{14}$ molecules) in each initial round of selection.^{34,41,42} The deoxyribozyme pool strand was 5'-CGAAGTCGCCATCTCTTC- N_x -ATAGTGAGTCGAT-TAAGCTGATCCTGATGG-3', where the two underlined regions denote the substrate binding arms. The 5'-CGAA was replaced with 5'-CC and the 3'-terminus was ...TATTA-3' for individual deoxyribozymes prepared by solid-phase synthesis and used in the single-turnover *in trans* kinetic assays. The two PCR primers used during selection were 5'-CGAAGTCGC-CATCTCTTC-3' (5'-phosphorylated to enable ligation by T4 RNA ligase) and 5'-(AAC)₄XCCATCAGGATCAGCTTAAT-ACGACTCACTAT-3' (where X denotes Glen Spacer 18). The RNA substrate used during selection was 5'-triphosphorylated 5'-ggaagagauggcgacgg-3', prepared by *in vitro* transcription using T7 RNA polymerase and a double stranded DNA template prepared by annealing two synthetic DNA oligonucleotides;⁴⁴ the 3'-terminus of the RNA substrate was joined by T4 RNA ligase to the 5'-terminus of the deoxyribozyme pool strand. The DNA-anchored tripeptide substrate was 5'-GGATAATACGACTCACTAT-3', joined via its 3'-terminus to a hexa(ethylene glycol) linker and the Cys-Tyr-Ala tripeptide through a disulfide bond as described.³⁴ The same DNA-anchored tripeptide substrate and 5'-triphosphorylated RNA substrate were used for single-turnover *in trans* kinetic assays of individual deoxyribozymes.

DNA Cleavage Assays. The DNA cleavage assays were performed under single-turnover *in trans* conditions with the procedure described previously.³⁸ The DNA substrate was 5'-³²P-radiolabeled using γ -³²P-ATP and T4 polynucleotide kinase. The final incubation conditions were 70 mM HEPES, pH 7.5, 1 mM $ZnCl_2$, 20 mM $MnCl_2$, 40 mM $MgCl_2$, and 150 mM NaCl at 37 °C, with 10 nM DNA substrate and 1 μ M deoxyribozyme (1:100) in 20 μ L total volume. The Mn^{2+} and Mg^{2+} were omitted in some assays as indicated. Samples were separated by 20% PAGE and quantified with a PhosphorImager.

Nucleopeptide Linkage Formation Assays. The nucleopeptide linkage formation assays were performed under single-turnover *in trans* conditions with the procedure described previously.³⁴ The DNA-anchored tripeptide substrate was 5'-³²P-radiolabeled using γ -³²P-ATP and T4 polynucleotide kinase. The final incubation conditions were 50 mM HEPES, pH 7.5, 20 mM $MnCl_2$, 40 mM $MgCl_2$, 150 mM NaCl, and 2 mM KCl at 37 °C, with 25 nM DNA-anchored tripeptide substrate, 0.5 μ M deoxyribozyme, and 1 μ M 5'-triphosphorylated RNA substrate (1:20:40) in 20 μ L total volume. Samples were separated by 20% PAGE and quantified with a PhosphorImager.

MALDI Mass Spectrometry. Samples for MALDI mass spectrometry were prepared using the procedure described previously.³⁰ DNA cleavage samples were prepared using 100 pmol of DNA substrate and 200 pmol of deoxyribozyme in 50 μ L reaction volume. Nucleopeptide linkage formation samples for mass spectrometry were prepared using 500 pmol of DNA-anchored tripeptide substrate, 600 pmol of deoxyribozyme, and 800 pmol of 5'-triphosphorylated RNA substrate in 50 μ L reaction volume. All mass spectra were obtained in the mass spectrometry laboratory of the UIUC School of Chemical Sciences.

■ ASSOCIATED CONTENT

📄 Supporting Information

Deoxyribozyme sequences, MALDI mass spectrometry data, and additional kinetic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Phone: 217-244-4489. Fax: 217-244-8024. E-mail: scott@scs.illinois.edu.

Author Contributions

[†]T.E.V. and J.S. are co-first authors of this manuscript.

Funding

This research was supported by grants to S.K.S. from the National Institutes of Health (R01GM065966), the Defense Threat Reduction Agency (HDTRA1-09-1-0011), and the National Science Foundation (CHE0842534).

Notes

The authors declare no competing financial interest.

■ REFERENCES

- (1) Roth, A.; Breaker, R. R. The Structural and Functional Diversity of Metabolite-Binding Riboswitches. *Annu. Rev. Biochem.* **2009**, *78*, 305–334.
- (2) Doudna, J. A.; Cech, T. R. The chemical repertoire of natural ribozymes. *Nature* **2002**, *418*, 222–228.
- (3) Leung, E. K.; Suslov, N.; Tuttle, N.; Sengupta, R.; Piccirilli, J. A. The mechanism of peptidyl transfer catalysis by the ribosome. *Annu. Rev. Biochem.* **2011**, *80*, 527–555.
- (4) Tuerk, C.; Gold, L. Systematic evolution of ligands by exponential enrichment: RNA ligands to bacteriophage T4 DNA polymerase. *Science* **1990**, *249*, 505–510.
- (5) Ellington, A. D.; Szostak, J. W. In vitro selection of RNA molecules that bind specific ligands. *Nature* **1990**, *346*, 818–822.
- (6) Robertson, D. L.; Joyce, G. F. Selection in vitro of an RNA enzyme that specifically cleaves single-stranded DNA. *Nature* **1990**, *344*, 467–468.
- (7) Ellington, A. D.; Szostak, J. W. Selection in vitro of single-stranded DNA molecules that fold into specific ligand-binding structures. *Nature* **1992**, *355*, 850–852.
- (8) Bock, L. C.; Griffin, L. C.; Latham, J. A.; Vermaas, E. H.; Toole, J. J. Selection of single-stranded DNA molecules that bind and inhibit human thrombin. *Nature* **1992**, *355*, 564–566.
- (9) Breaker, R. R.; Joyce, G. F. A DNA enzyme that cleaves RNA. *Chem. Biol.* **1994**, *1*, 223–229.
- (10) Joyce, G. F. Directed Evolution of Nucleic Acid Enzymes. *Annu. Rev. Biochem.* **2004**, *73*, 791–836.
- (11) Joyce, G. F. Forty Years of In Vitro Evolution. *Angew. Chem., Int. Ed.* **2007**, *46*, 6420–6436.
- (12) Silverman, S. K. Artificial Functional Nucleic Acids: Aptamers, Ribozymes, and Deoxyribozymes Identified by In Vitro Selection. In *Functional Nucleic Acids for Analytical Applications*; Li, Y., Lu, Y., Eds.;

Springer Science + Business Media, LLC: New York, 2009; pp 47–108.

(13) Silverman, S. K. Catalytic DNA (deoxyribozymes) for synthetic applications—current abilities and future prospects. *Chem. Commun.* **2008**, 3467–3485.

(14) Schlosser, K.; Li, Y. Biologically inspired synthetic enzymes made from DNA. *Chem. Biol.* **2009**, *16*, 311–322.

(15) Silverman, S. K. DNA as a Versatile Chemical Component for Catalysis, Encoding, and Stereocontrol. *Angew. Chem., Int. Ed.* **2010**, *49*, 7180–7201.

(16) Silverman, S. K. In Vitro Selection and Application of Nucleic Acid Enzymes (Ribozymes and Deoxyribozymes). In *Wiley Encyclopedia of Chemical Biology*; Begley, T. P., Ed.; John Wiley and Sons: Hoboken, NJ, 2009; DOI: 10.1002/9780470048672.wecb406.

(17) Li, Y.; Geyer, C. R.; Sen, D. Recognition of anionic porphyrins by DNA aptamers. *Biochemistry* **1996**, *35*, 6911–6922.

(18) Sabeti, P. C.; Unrau, P. J.; Bartel, D. P. Accessing rare activities from random RNA sequences: the importance of the length of molecules in the starting pool. *Chem. Biol.* **1997**, *4*, 767–774.

(19) Irvine, D.; Tuerk, C.; Gold, L. SELEXION. Systematic evolution of ligands by exponential enrichment with integrated optimization by non-linear analysis. *J. Mol. Biol.* **1991**, *222*, 739–761.

(20) Knight, R.; De Sterck, H.; Markel, R.; Smit, S.; Oshmyansky, A.; Yarus, M. Abundance of correctly folded RNA motifs in sequence space, calculated on computational grids. *Nucleic Acids Res.* **2005**, *33*, 5924–5935.

(21) Sun, F.; Galas, D.; Waterman, M. S. A mathematical analysis of in vitro molecular selection-amplification. *J. Mol. Biol.* **1996**, *258*, 650–660.

(22) Gevertz, J.; Gan, H. H.; Schlick, T. In vitro RNA random pools are not structurally diverse: a computational analysis. *RNA* **2005**, *11*, 853–863.

(23) Levine, H. A.; Nilsen-Hamilton, M. A mathematical analysis of SELEX. *Comput. Biol. Chem.* **2007**, *31*, 11–35.

(24) Chushak, Y.; Stone, M. O. In silico selection of RNA aptamers. *Nucleic Acids Res.* **2009**, *37*, e87.

(25) Kim, N.; Izzo, J. A.; Elmetwaly, S.; Gan, H. H.; Schlick, T. Computational generation and screening of RNA motifs in large nucleotide sequence pools. *Nucleic Acids Res.* **2010**, *38*, e139.

(26) Huang, F.; Bugg, C. W.; Yarus, M. RNA-Catalyzed CoA, NAD, and FAD synthesis from phosphopantetheine, NMN, and FMN. *Biochemistry* **2000**, *39*, 15548–15555.

(27) Coleman, T. M.; Huang, F. RNA-catalyzed thioester synthesis. *Chem. Biol.* **2002**, *9*, 1227–1236.

(28) Coleman, T. M.; Huang, F. Optimal Random Libraries for the Isolation of Catalytic RNA. *RNA Biol.* **2005**, *2*, 129–136.

(29) Legiewicz, M.; Lozupone, C.; Knight, R.; Yarus, M. Size, constant sequences, and optimal selection. *RNA* **2005**, *11*, 1701–1709.

(30) Chandra, M.; Sachdeva, A.; Silverman, S. K. DNA-catalyzed sequence-specific hydrolysis of DNA. *Nat. Chem. Biol.* **2009**, *5*, 718–720.

(31) Xiao, Y.; Wehrmann, R. J.; Ibrahim, N. A.; Silverman, S. K. Establishing Broad Generality of DNA Catalysts for Site-Specific Hydrolysis of Single-Stranded DNA. *Nucleic Acids Res.* **2012**, *40*, 1778–1786.

(32) Pradeepkumar, P. I.; Höbartner, C.; Baum, D. A.; Silverman, S. K. DNA-Catalyzed Formation of Nucleopeptide Linkages. *Angew. Chem., Int. Ed.* **2008**, *47*, 1753–1757.

(33) Sachdeva, A.; Silverman, S. K. DNA-Catalyzed Serine Side Chain Reactivity and Selectivity. *Chem. Commun.* **2010**, *46*, 2215–2217.

(34) Wong, O. Y.; Pradeepkumar, P. I.; Silverman, S. K. DNA-Catalyzed Covalent Modification of Amino Acid Side Chains in Tethered and Free Peptide Substrates. *Biochemistry* **2011**, *50*, 4741–4749.

(35) Wong, O. Y.; Mulcrone, A. E.; Silverman, S. K. DNA-Catalyzed Reductive Amination. *Angew. Chem., Int. Ed.* **2011**, *50*, 11679–11684.

- (36) Sachdeva, A.; Silverman, S. K. DNA-catalyzed reactivity of a phosphoramidate functional group and formation of an unusual pyrophosphoramidate linkage. *Org. Biomol. Chem.* **2012**, *10*, 122–125.
- (37) Sachdeva, A.; Chandra, M.; Chandrasekar, J.; Silverman, S. K. Covalent Tagging of Phosphorylated Peptides By Phosphate-Specific Deoxyribozymes. *ChemBioChem* **2012**, *13*, 654–657.
- (38) Xiao, Y.; Chandra, M.; Silverman, S. K. Functional Compromises among pH Tolerance, Site Specificity, and Sequence Tolerance for a DNA-Hydrolyzing Deoxyribozyme. *Biochemistry* **2010**, *49*, 9630–9637.
- (39) Xiao, Y.; Allen, E. C.; Silverman, S. K. Merely two mutations switch a DNA-hydrolyzing deoxyribozyme from heterobimetallic (Zn^{2+}/Mn^{2+}) to monometallic (Zn^{2+} -only) behavior. *Chem. Commun.* **2011**, *47*, 1749–1751.
- (40) Dokukin, V.; Silverman, S. K. Lanthanide Ions as Required Cofactors for DNA Catalysts. *Chem. Sci.* **2012**, *3*, 1707–1714.
- (41) Flynn-Charlebois, A.; Wang, Y.; Prior, T. K.; Rashid, I.; Hoadley, K. A.; Coppins, R. L.; Wolf, A. C.; Silverman, S. K. Deoxyribozymes with 2'-5' RNA Ligase Activity. *J. Am. Chem. Soc.* **2003**, *125*, 2444–2454.
- (42) Kost, D. M.; Gerdt, J. P.; Pradeepkumar, P. I.; Silverman, S. K. Controlling regioselectivity and site-selectivity in RNA ligation by Zn^{2+} -dependent deoxyribozymes that use 2',3'-cyclic phosphate RNA substrates. *Org. Biomol. Chem.* **2008**, *6*, 4391–4398.
- (43) Sheppard, T. L.; Ordoukhanian, P.; Joyce, G. F. A DNA enzyme with *N*-glycosylase activity. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 7802–7807.
- (44) Milligan, J. F.; Groebe, D. R.; Witherell, G. W.; Uhlenbeck, O. C. Oligoribonucleotide synthesis using T7 RNA polymerase and synthetic DNA templates. *Nucleic Acids Res.* **1987**, *15*, 8783–8798.